

Comments, suggestions, corrections, and references welcomed!

SOME RESULTS ON CAPABLE GROUPS OF PRIME EXPONENT AND CLASS TWO

ARTURO MAGIDIN

ABSTRACT. This note collects several results on the capability of p -groups of class two and prime exponent. Among our results, we settle the 4-generator case for this class.

INTRODUCTION.

In this note we collect sundry results related to the capability of finite p -groups of class two and prime exponent, using the approach introduced in [6]. We restrict to odd primes, since the case of abelian groups is well understood. I owe a considerable amount of the material here to discussions with David McKinnon, who helped me clarify much of the background and ideas.

A group G is said to be *capable* if and only if there exists a group K such that $G = K/Z(K)$. For p -groups (groups of finite prime-power order) capability is closely related to their classification. Baer characterized the capable groups which are direct sums of cyclic groups in [2]; the capable extra-special p -groups were characterized by Beyl, Felgner, and Schmid in [3] (only the dihedral group of order 8 and the extra-special groups of order p^3 and exponent p are capable); the three authors also describe the metacyclic groups which are capable. The author characterized the 2-generated capable p -groups of class two [5, 7] (for odd p , independently obtained in part by Bacon and Kappe in [1]).

In [6] we analyzed the special case of p -groups of class two and exponent p , where p is an odd prime. We described a way to translate the problem into a statement of linear algebra, and obtained several results. We will introduce what I hope is clearer notation here, reprove some of the results using it, and we will also be able to obtain new results. Some results from [6] we will simply quote, and we direct the reader there for any omitted proofs.

1. THE SET-UP.

We begin with a general construction and some observations.

Definition 1.1. Let V_1 and V_2 be vector spaces, and let $\{\ell_i: V_1 \rightarrow V_2\}_{i \in I}$ be a nonempty family of linear transformations. For every subspace X of V_1 , we define $X^* < V_2$ to be:

$$X^* = \langle \ell_i(X) \mid i \in I \rangle.$$

For every subspace Y of V_2 , we define $Y^* < V_1$ to be:

$$Y^* = \bigcap_{i \in I} \ell_i^{-1}(Y).$$

2000 *Mathematics Subject Classification.* Primary 20D15, Secondary 20F12.

Note that for all subspaces of V_1 , if $X \subset X'$, then $X^* \subset X'^*$; likewise, for subspaces of V_2 , if $Y \subset Y'$, then $Y^* \subset Y'^*$.

Theorem 1.2. *Let V_1 and V_2 be vector spaces, and let $\{\ell_i: V_1 \rightarrow V_2\}_{i \in I}$ be a nonempty family of linear transformations. The operator on the subspaces of V_1 defined by $X \mapsto X^{**}$ is a closure operator; that is, it is increasing, isotone, and idempotent. Moreover, if X is a subspace of V_1 , then $X^* = X^{***}$.*

Proof. It is clear that $X \subset X^{**}$, so the operator is increasing; since $X \subset X'$ implies $X^* \subset X'^*$, which in turn implies $X^{**} \subset X'^{**}$, the operator is isotone.

To show the operator is idempotent, we want to show that $X^{**} = (X^{**})^{**}$. For simplicity, write $X^{**} = Z$. Since the operator is increasing, we know that $Z \subset Z^{**}$. By construction, we also have that $\ell_i(Z) \subset X^*$ for each i , so $Z^* \subset X^*$. From this, we get that $Z^{**} \subset X^{**} = Z$, as desired.

Finally, let $X \subset V_1$. Since $X \subset X^{**}$, we must have $X^* \subset X^{***}$. Conversely, from the definition of X^{***} it follows that $\ell_i(X^{**}) \subset X^*$ for all i , so $X^{***} \subset X^*$, giving equality. $\square$

The dual property is true for subspaces of V_2 :

Theorem 1.3. *Let V_1 and V_2 be vector spaces, and let $\{\ell_i: V_1 \rightarrow V_2\}_{i \in I}$ be a nonempty family of linear transformations. The operator on subspaces of V_2 defined by $Y \mapsto Y^{**}$ is an interior operator; that is, it is decreasing, isotone, and idempotent. Moreover, if Y is a subspace of V_2 , then $Y^* = Y^{***}$.*

Proof. As before, the operator is isotone. Since $Y^{**} = \langle \ell_i(Y^*) \rangle$, and $Y^* \subset \ell_i^{-1}(Y)$ for each i , it follows that $Y^{**} \subset Y$, so the operator is decreasing.

To show the operator is idempotent, set $Z = Y^{**}$. Since $Y^* \subset Y^{***}$, we must have $Z = Y^{**} \subset Y^{****} = Z^{**}$. Since the other inclusion always holds, this shows that $Z = Z^{**}$, as desired.

Finally, since $Y^{**} \subset Y$, it follows that $Y^{***} \subset Y^*$. But since the operator $**$ on subspaces of V_1 is increasing, we also know that $Y^* \subset Y^{***}$, giving equality. $\square$

Definition 1.4. Let V_1, V_2 be vector spaces, and let $\{\ell_i: V_1 \rightarrow V_2\}_{i \in I}$ be a family of linear transformations. We will say that a subspace X of V_1 is $\{\varphi_i\}_{i \in I}$ -closed (or simply *closed* if there is no danger of ambiguity) if and only if $X = X^{**}$.

Recall that if V is a vector space and k a positive integer, the Grassmannian $Gr(k, V)$ is defined to be the set of all k -dimensional subspaces of V . It has more algebraic structure than simply being a set, but for now we will simply use it as convenient notation.

We fix an odd prime p throughout. Let n be an integer greater than 1. We define three vector spaces with distinguished bases:

Definition 1.5. Let $U(n)$ be the vector space over the finite field $\mathbb{F}_p$ of p elements, with basis vectors $x_1, \dots, x_n$. Let $V(n)$ be the vector space over $\mathbb{F}_p$ of dimension $\binom{n}{2}$, with basis vectors v_{ji} , $1 \leq i < j \leq n$. Let $W(n)$ be the vector space over $\mathbb{F}_p$ of dimension $2\binom{n}{2} + 2\binom{n}{3}$, with basis vectors w_{jik} , $1 \leq i < j \leq n, i \leq k \leq n$.

If there is no danger of ambiguity, and n is understood from context, we will simply use U, V , and W instead of $U(n), V(n)$, and $W(n)$. This will be the case in most of our applications.

We define two families of n linear operators:

Definition 1.6. The linear maps $\psi_i: U \rightarrow V$, $i = 1, \dots, n$ are defined to be:

$$\psi_i(x_j) = \begin{cases} v_{ji} & \text{if } i < j \\ -v_{ij} & \text{if } i > j \\ \mathbf{0} & \text{if } i = j. \end{cases}$$

Definition 1.7. The linear maps $\varphi_k: V \rightarrow W$, $k = 1, \dots, n$, are defined to be:

$$\varphi_k(v_{ji}) = \begin{cases} w_{jik} & \text{if } k \geq i, \\ w_{jki} - w_{ikj} & \text{if } k < i. \end{cases}$$

Note that there is a natural identification of V with $U \wedge U$, and of W with $(U \wedge U) \otimes U$ modulo the Jacobi identity:

$$(u_i \wedge u_j) \otimes u_k + (u_j \wedge u_k) \otimes u_i + (u_k \wedge u_i) \otimes u_j = 0.$$

Under these identifications, the maps ψ_i correspond to $\psi_i(\mathbf{u}) = \mathbf{u} \wedge u_i$, and the maps φ_k correspond to $\varphi_k(\mathbf{v} \wedge \mathbf{w}) = (\mathbf{v} \wedge \mathbf{w}) \otimes u_k$.

The definitions of U , V , W , ψ_j , and φ_k are derived from a group theoretic construction. Namely: let F be the free product of n cyclic groups of order p , generated by $x_1, \dots, x_n$ respectively. Let F_k be the k -th term of the lower central series of F . Then U corresponds to $F^{\text{ab}} = F/F_2$; V corresponds to F_2/F_3 , by identifying $[x_j, x_i]$ with v_{ji} ; and W corresponds to F_3/F_4 , identifying $[[x_j, x_i], x_k]$ with w_{jik} . The linear maps ψ_i and φ_i correspond to the maps induced by the commutator bracket $[-, x_j]$ from F/F_2 to F_2/F_3 , and from F_2/F_3 to F_3/F_4 , respectively. We could collect all of this information by making $U \oplus V \oplus W$ into a Lie algebra, via the Lie product given by the commutator bracket, but we will not go through the details here since we will not use the Lie algebra structure.

The connection to capability is explored and explained in [6]; we give a very brief summary here: let G be any p -group of class two and exponent p , with G^{ab} of rank n . Let $g_1, \dots, g_n$ be elements of G that project onto a basis of G^{ab} , and let $K = F/F_3$. Then K is the relatively free group of rank n in the variety of nilpotent groups of class at most two and exponent p , so there is a unique homomorphism $K \rightarrow G$ mapping x_i to g_i . Let N be the kernel. Then $N \subset [K, K]$. Thus, N corresponds to a subspace X of V (since $[K, K] = F_2/F_3$ corresponds to V). We then have:

Theorem 1.8 (Theorem 3.6 in [6]). *For $n > 1$, and let G , K , and N be as in the previous paragraph; identify $[K, K]$ with V by letting $[x_j, x_i]$ correspond to v_{ji} , $1 \leq i < j \leq n$, and let X be the subspace of V corresponding to N . Then G is capable if and only if X is closed with respect to the φ_i .*

2. THE LINEAR ALGEBRA.

We collect here some observations on the spaces V and W , and the linear transformations φ_k defined above.

Definition 2.1. Let i, j be integers, $1 \leq i < j \leq n$. We let $\pi_{ji}: V \rightarrow \langle v_{ji} \rangle$ be the canonical projection.

Definition 2.2. Let i, j, k be integers, $1 \leq i < j \leq n$, $i \leq k \leq n$. We let $\pi_{jik}: W \rightarrow \langle w_{jik} \rangle$ be the canonical projection.

Definition 2.3. Let i be an integer, $1 \leq i \leq n$. We let

$$\Pi_i: V \rightarrow \langle v_{i1}, v_{i2}, \dots, v_{i(i-1)}, v_{(i+1)i}, \dots, v_{ni} \rangle$$

be the canonical projection.

Lemma 2.4. For each k , φ_k is one-to-one.

Lemma 2.5. Let $\mathbf{w} \in \varphi_i(V)$. If $\pi_{rst}(\mathbf{w}) \neq \mathbf{0}$, then $s \leq i \leq t$; moreover, exactly one of the inequalities is strict.

Lemma 2.6. Fix n, i, j , with $n > 1$ and $1 \leq i < j \leq n$. Then $\varphi_i(V) \cap \varphi_j(V) = \{\mathbf{0}\}$.

Proof. Assume $\varphi_i(\mathbf{v}) \in \varphi_j(V)$, and that $\pi_{sr}(\mathbf{v}) \neq \mathbf{0}$. If $r \leq i$, then $\pi_{sri}(\varphi_i(\mathbf{v})) \neq \mathbf{0}$; but since $\varphi_i(\mathbf{v}) \in \varphi_j(V)$, this implies that $r \leq j \leq i$, which is impossible. If, on the other hand, $i < r$, then $\pi_{sir}(\varphi_i(\mathbf{v})) \neq \mathbf{0}$ and $\pi_{ris}(\varphi_i(\mathbf{v})) \neq \mathbf{0}$; again, since $\varphi_i(\mathbf{v}) \in \varphi_j(V)$, Lemma 2.5 implies that $s = r = j$, which is also impossible. $\square$

Lemma 2.7. Fix $n > 1$, $r \leq n$. Let $i_1, \dots, i_r$ be integers, $1 \leq i_1 \leq \dots \leq i_r \leq n$. Then

$$\varphi_{i_1}(V) \cap \langle \varphi_{i_2}(V), \dots, \varphi_{i_r}(V) \rangle$$

is of dimension $\binom{r-1}{2}$, with basis given by all vectors $w_{ai_1b} - w_{bi_1a}$, $a, b \in \{i_2, \dots, i_r\}$, $a > b$. A basis for the pullback

$$\varphi_{i_1}^{-1}(\langle \varphi_{i_2}(V), \dots, \varphi_{i_r}(V) \rangle)$$

is given by $\{v_{ab}\}$, with $a, b \in \{i_2, \dots, i_r\}$, $a > b$.

Proof. The result is true for $r = 1$ and $r = 2$. Assume then that $r \geq 3$. Note that the vectors $w_{ai_1b} - w_{bi_1a}$ are indeed in the intersection; for any choice of $a > b$ as given, we have $\varphi_{i_1}(v_{ab}) = w_{ai_1b} - w_{bi_1a} = \varphi_b(v_{ai_1}) - \varphi_a(v_{bi_1})$.

Let $S = \langle \varphi_{i_2}(V), \dots, \varphi_{i_r}(V) \rangle$. Assume that $\varphi_{i_1}(\mathbf{v}) \in S$, and that $\pi_{ts}(\mathbf{v}) \neq \mathbf{0}$, where $1 \leq s < t \leq n$. Since $\varphi_{i_1}(\mathbf{v}) \in S$, and every vector in S has trivial image under all π_{abi_1} , it follows that $s > i_1$. Therefore $\varphi_{i_1}(v_{ts}) = w_{ti_1s} - w_{si_1t}$, and so we have:

$$\varphi_{i_1}(\mathbf{v}) = \varphi_{i_1} \left(\sum_{i_1 < s < t \leq n} \alpha_{ts} v_{ts} \right) = \sum_{i_1 < s < t \leq n} \alpha_{ts} (w_{ti_1s} - w_{si_1t}).$$

Let $\mathbf{w} \in S$; if $\pi_{jik}(\mathbf{w}) \neq \mathbf{0}$ with $i < i_2$, then we must have $k = i_c$ for some $c \in \{2, \dots, r\}$. Therefore, if $\varphi_{i_1}(\mathbf{v}) \in S$, and $\pi_{ts}(\mathbf{v}) = \alpha_{ts} v_{ts} \neq \mathbf{0}$, then s and t are in $\{i_2, \dots, i_r\}$. Thus we may rewrite $\varphi_{i_1}(\mathbf{v})$ as:

$$\varphi_{i_1}(\mathbf{v}) = \sum_{2 \leq s < t \leq r} \alpha_{ts} (w_{ti_1i_s} - w_{si_1i_t}) = \sum_{2 \leq s < t \leq r} \varphi_{i_s}(\alpha_{ts} v_{ti_1}) - \varphi_{i_t}(v_{si_1}).$$

This proves the lemma. $\square$

Corollary 2.8. Fix $n > 1$, $r \leq n$, and let $1 \leq i_1 \leq \dots \leq i_r \leq n$ be integers. Then

$$\dim(\langle \varphi_{i_1}(V), \dots, \varphi_{i_r}(V) \rangle) = r \binom{n}{2} - \binom{r}{3}.$$

Proof. Each φ_k is injective, so we have:

$$\begin{aligned} & \dim(\langle \varphi_{i_1}(V), \dots, \varphi_{i_r}(V) \rangle) \\ &= \left(\sum_{k=1}^r \dim(\varphi_{i_k}(V)) \right) - \left(\sum_{k=1}^{r-2} \dim(\varphi_{i_{r-k-1}}(V) \cap \langle \varphi_{i_r}(V), \dots, \varphi_{i_{r-k}}(V) \rangle) \right) \\ &= r \binom{n}{2} - \left(\sum_{k=1}^r \binom{k-1}{2} \right) = r \binom{n}{2} - \binom{r}{3}, \end{aligned}$$

as claimed. $\square$

We have concentrated on the “lowest index” for simplicity. Of course, given the definitions, our treatment has symmetry; for example:

Proposition 2.9. *Let $i_1, \dots, i_r \in \{1, \dots, n\}$ be pairwise distinct. Then*

$$\varphi_{i_1}(V) \cap \langle \varphi_{i_2}(V), \dots, \varphi_{i_r}(V) \rangle$$

has dimension $\binom{r-1}{2}$. Moreover, a basis for the pullback is given by the vectors v_{ab} , with $a, b \in \{i_2, \dots, i_r\}$, and $a > b$.

Proof. It is easy to verify that

$$\varphi_{i_1}(V) \cap \langle \varphi_{i_2}(V), \dots, \varphi_{i_r}(V) \rangle$$

is generated by the vectors w_{abi_1} for $a, b \in \{i_2, \dots, i_r\}$, $a, i_1 > b$, and the vectors $w_{ai_1b} - w_{bi_1a}$ when $b > i_1$. These vectors are linearly independent, and pulling them back gives the desired result. $\square$

As we will see below, these descriptions allow us to place bounds on $\dim(X^*)$ in terms of $\dim(X)$ and n .

3. DIMENSION COUNTING.

In this section we will prove some bounds on $\dim(X^*)$ in terms of $\dim(X)$ and n . These bounds will give sufficient conditions for X to be closed in terms of its dimension.

Lemma 3.1 (Prop. 4.6 in [6]). *Fix $n > 1$, and let X be a subspace of V . If $\dim(X) = 1$, then $\dim(X^*) = n$; if $\dim(X) = 2$, then $\dim(X^*) = 2n$.*

In [6] we proved this by considering the collection of images of a basis of X under $\varphi_1, \dots, \varphi_n$, and showing they would necessarily be linearly independent. We give a new proof here:

Proof. The results are trivial if $n = 2$, so assume $n \geq 3$. We will prove the contrapositive. Note that $\dim(X^*) \leq n \dim(X)$ always holds, so assume that $\dim(X^*) < n \dim(X)$. We will show that $\dim(X) \geq 3$.

Let k be any integer such that

$$\varphi_k(X) \cap \langle \varphi_{k+1}(X), \dots, \varphi_n(X) \rangle \neq \{0\}.$$

Such a k must exist, for if all such intersections were trivial then we would have $\dim(X^*) = n \dim(X)$. Let

$$\mathbf{v} \in \varphi_k^{-1}(\langle \varphi_{k+1}(X), \dots, \varphi_n(X) \rangle) - \{0\}.$$

Then from Lemma 2.7 and the injectivity of φ_k , we deduce that we can write

$$\mathbf{v} = \sum_{k+1 \leq i < j \leq n} \alpha_{ji} v_{ji},$$

and not all α_{ji} are zero. Therefore,

$$\begin{aligned} \varphi_k(\mathbf{v}) &= \sum_{k+1 \leq i < j \leq n} \alpha_{ji} (w_{jki} - w_{ikj}) \\ &= \sum_{k+1 \leq i < j \leq n} \alpha_{ji} (\varphi_i(v_{jk}) - \varphi_j(v_{ik})) \\ &= \sum_{i=k+1}^n \varphi_i \left(\sum_{j=i+1}^n \alpha_{ji} v_{jk} - \sum_{j=k+1}^{i-1} \alpha_{ij} v_{jk} \right). \end{aligned}$$

By injectivity of the φ and the choice of $\mathbf{v}$, we again conclude that for each value of i , $i = k+1, \dots, n$,

$$\mathbf{v}_i = \left(\sum_{j=i+1}^n \alpha_{ji} v_{jk} - \sum_{j=k+1}^{i-1} \alpha_{ij} v_{jk} \right) \in X.$$

For simplicity, set $\alpha_{ij} = -\alpha_{ji}$, and $\alpha_{ii} = 0$. Then we can rewrite $\mathbf{v}_i$ as:

$$\mathbf{v}_i = \sum_{k+1 \leq j \leq n} \alpha_{ji} v_{jk}.$$

Let $S = \{\mathbf{v}_{k+1}, \mathbf{v}_{k+2}, \dots, \mathbf{v}_n\}$. Note that if a subset $T \subset S$ is linearly independent, then so is $T \cup \{\mathbf{v}\}$. We will show that there are at least two vectors in S that are not multiples of each other; this will show that the dimension of X is at least three, proving the result.

Indeed, fix i_0 and j_0 such that $\alpha_{j_0 i_0} \neq 0$, $k < i_0 < j_0 \leq n$. We have that

$$\mathbf{v}_{i_0} = \sum_{j=k+1}^n \alpha_{ji_0} v_{jk} \quad \text{and} \quad \mathbf{v}_{j_0} = \sum_{j=k+1}^n \alpha_{jj_0} v_{jk},$$

so $\pi_{j_0 k}(\mathbf{v}_{i_0}) \neq \mathbf{0}$ and $\pi_{i_0 k}(\mathbf{v}_{i_0}) = \mathbf{0}$; on the other hand, $\pi_{i_0 k}(\mathbf{v}_{j_0}) \neq \mathbf{0}$ and $\pi_{j_0 k}(\mathbf{v}_{j_0}) = \mathbf{0}$. So neither of $\mathbf{v}_{i_0}$ and $\mathbf{v}_{j_0}$ is a scalar multiple of the other, and $\{\mathbf{v}_{i_0}, \mathbf{v}_{j_0}, \mathbf{v}\}$ is a linearly independent subset of X . Therefore, the dimension of X must be at least 3, as claimed. $\square$

With similar arguments we can obtain a (naive) lower bound for the dimension of X^* in terms of $\dim(X)$. We have:

Lemma 3.2. *Fix $n \geq 3$, and let X be a subspace of V , with $\dim(X) = k$. Then*

$$nk \geq \dim(X^*) \geq \begin{cases} nk - \left(\frac{k-1}{2}\right)^2 & \text{if } k \text{ is odd,} \\ nk - \binom{k}{2} \left(\frac{k-2}{2}\right) & \text{if } k \text{ is even.} \end{cases}$$

Proof. The upper bound is trivial. Let

$$Z_i = \varphi_i(X) \cap \langle \varphi_{i+1}(X), \dots, \varphi_n(X) \rangle$$

for $i = 1, 2, \dots, n-2$. Note that

$$\dim(X^*) = nk - \sum_{i=1}^{n-2} \dim(Z_i).$$

Let $1 \leq i_1 < i_2 < \dots < i_r \leq n-2$ be the indices for which Z_{i_j} is nontrivial. The dimension of Z_{i_1} is at most $k-2$; this since, as above, if $\mathbf{v}$ is in $\varphi_{i_1}^{-1}(Z_{i_1})$, then $\Pi_k(\mathbf{v}) \neq \mathbf{0}$ implies $k \in \{i_2, \dots, i_r\}$; and we need at least two linearly independent vectors with nontrivial image under Π_{i_1} to account for the intersection. Likewise, the dimension of Z_{i_2} is at most $k-4$, since the two vectors with nontrivial image under Π_{i_1} cannot be involved in this intersection. Continuing this way, we obtain

$$\sum_{j=1}^r \dim(Z_{i_j}) \leq (k-2) + (k-4) + \dots + (k-2r).$$

Moreover, note that $2r < k$. If k is odd, say $k = 2\ell + 1$, then this gives

$$\begin{aligned} \sum_{j=1}^r \dim(Z_{i_j}) &\leq (k-2) + (k-4) + \dots + (k-2\ell) \\ &= \ell k - 2(1 + 2 + \dots + \ell) \\ &= 2\ell^2 + \ell - \ell^2 - \ell \\ &= \left(\frac{k-1}{2}\right)^2. \end{aligned}$$

On the other hand, if k is even, $k = 2\ell$, then

$$\begin{aligned} \sum_{j=1}^r \dim(Z_{i_j}) &\leq (k-2) + (k-4) + \dots + (k-2(\ell-1)) \\ &= (\ell-1)k - 2(1 + 2 + \dots + (\ell-1)) \\ &= 2\ell^2 - 2\ell - \ell^2 + \ell \\ &= \left(\frac{k}{2}\right)^2 - \frac{k}{2} \\ &= \left(\frac{k}{2}\right) \left(\frac{k-2}{2}\right). \end{aligned}$$

This proves the lower bound. $\square$

Remark 3.3. We can improve upon the lower bound with more care; for example, as we will see below, whereas the lemma only gives $\dim(X^*) \geq 16$ in the case $n = 4$ and $k = 5$, we can actually prove that we will have $\dim(X^*) \geq 17$.

Though very rough, the lower bounds established below give us a sufficient condition for X to be closed in terms only of $\dim(X)$ and n :

Theorem 3.4. *Let $n > 1$, and let X be a subspace of V . If $\dim(X) = k$, then X is closed whenever $k < 2\sqrt{n}$.*

Proof. Let X' be a subspace of dimension $k + 1$; we show that if k satisfies the given inequality then $\dim(X'^*) > \dim(X^*)$.

From Lemma 3.2 we know that $\dim(X^*) \leq nk$ and

$$\dim(X'^*) \geq \begin{cases} n(k+1) - \left(\frac{k+1}{2}\right)\left(\frac{k-1}{2}\right) & \text{if } k \text{ is odd,} \\ n(k+1) - \left(\frac{k+1}{2}\right)^2 & \text{if } k \text{ is even.} \end{cases}$$

Thus, $\dim(X^*) < \dim(X'^*)$ will certainly hold provided that

$$\begin{aligned} k^2 &< 4n + 1 && \text{if } k \text{ is odd;} \\ k^2 &< 4n && \text{if } k \text{ is even.} \end{aligned}$$

It is easy to verify that these inequalities hold if and only if $k < 2\sqrt{n}$, regardless of the parity of k . This proves that $\dim(X^*) < \dim(X'^*)$ for any X' of dimension $k + 1$, provided $k < 2\sqrt{n}$.

Since any X' that satisfies $X \subseteq X' \subseteq X^{**}$ must have $X'^* = X^*$, it follows that there are no subspaces of dimension $k + 1$ that lie between X and X^{**} ; this means that $X^{**} = X$, so X is closed, as claimed. $\square$

Translating into the group theoretic setting, we have:

Theorem 3.5. *Let G be a p -group of class two and exponent p , $\text{rank}(G^{\text{ab}}) = n$, and $\text{rank}([G, G]) = \ell$. If $\ell > \binom{n}{2} - 2\sqrt{n}$ then G is capable.*

We also have:

Corollary 3.6 (cf. Cor. 3.7 in [6]). *Let $n \geq 3$, and X be a subspace of V . If $\dim(X) \leq 4$, then X is closed.*

Proof. Let $\dim(X) = k$. Theorem 3.4 gives the result if $k \leq 3$, or if $k = 4 < n$. So we may assume $\dim(X) = n = 4$. From Lemma 3.2, we know that if $\dim(X') = 5$, then

$$16 \leq \dim(X'^*) \leq 20.$$

We claim that in fact we can sharpen the lower bound to $\dim(X'^*) \geq 17$. If so this will prove the proposition via the same argument as in the final paragraph of the proof of Theorem 3.4. There are only two possible values of i for which

$$Z'_i = \varphi_i(X') \cap \langle \varphi_{i+1}(X'), \dots, \varphi_n(X') \rangle$$

will be nontrivial: $i = 1$ and $i = 2$. The dimension of Z'_2 is at most 1, and the dimension of Z'_1 is at most 3; but the only way for Z'_1 to be of dimension 3 is if X' contains all of v_{32} , v_{42} , and v_{43} , in which case it is trivial to see that we must also have v_{21} , v_{31} , and v_{41} in X' . Since $\dim(X') = 5$, this is impossible. Thus, in fact, $\dim(Z'_2) \leq 1$ and $\dim(Z'_1) \leq 2$, so $\dim(X'^*) \geq 20 - 3 = 17$. This proves the claim, and the proposition. $\square$

4. SOME RESULTS.

We quote the following two results without proof:

Theorem 4.1 (Lemma 5.3 in [6]). *If X is a coordinate subspace (that is, it is generated by a subset of the v_{ji}) then X is closed.*

Theorem 4.2 (Lemma 5.7 in [6]). *Let m be fixed, $1 < m < n$, and assume that X_1 is a subspace of $\langle v_{ji} \mid 1 \leq i < j \leq m \rangle$, and X_2 is a subspace of $\langle v_{ji} \mid m < i < j \leq n \rangle$. Then $X_1 \oplus X_2$ is closed.*

In [6, Lemma 5.9] we proved a special case of the following result:

Lemma 4.3. *Let $n > 1$ be an integer, let I be a proper nonempty subset of $\{1, \dots, n\}$, and let $J = \{1, \dots, n\} - I$. Let*

$$\begin{aligned} V_I &= \langle v_{ji} \mid 1 \leq i < j \leq n; i, j \in I \rangle \\ V_J &= \langle v_{ji} \mid 1 \leq i < j \leq n; i, j \in J \rangle \\ V_{(I,J)} &= \langle v_{ji} \mid 1 \leq i < j \leq n; \text{ exactly one of } i, j \text{ is in } I, \text{ one in } J \rangle. \end{aligned}$$

Let X_I be a subspace of V_I , X_J be a subspace of V_J , and let

$$X = X_I \oplus X_J \oplus V_{(I,J)}.$$

Then

$$X^{**} = \left(\bigcap_{i \in I} \varphi_i^{-1} \left(\langle \varphi_k(X_I) \mid k \in I \rangle \right) \right) \oplus \left(\bigcap_{j \in J} \varphi_j^{-1} \left(\langle \varphi_k(X_J) \mid k \in J \rangle \right) \right) \oplus V_{(I,J)}.$$

In particular, X is $\{\varphi_k\}_{k=1}^n$ -closed if and only if X_I is $\{\varphi_i\}_{i \in I}$ -closed and X_J is $\{\varphi_j\}_{j \in J}$ -closed.

Proof. Let W' be the subspace of W generated by all w_{jik} in which exactly one or two of i, j, k are in I . First, we claim that $W' \subset X^*$. Suppose that exactly one of i, j, k lies in I . If $j \in I$, then $v_{ji} \in V_{(I,J)}$, so $\varphi_k(v_{ji}) = w_{jik} \in X^*$. If $i \in I$, then again $v_{ji} \in V_{(I,J)}$, so again we conclude that $w_{jik} \in X^*$. Finally, if $k \in I$, then either v_{jk} or v_{kj} lies in $V_{(I,J)}$ (whichever is appropriate); the image under φ_i of this vector is $\pm(w_{jik} - w_{kij})$. Since $w_{kij} \in X^*$ by the argument above it follows that $w_{jik} \in X^*$ as desired. The case when exactly two of i, j, k lie in I follows by symmetry, since exactly one of them will lie in J .

Next, we claim that

$$X^* = \langle \varphi_i(X_I) \mid i \in I \rangle \oplus \langle \varphi_j(X_J) \mid j \in J \rangle \oplus W'.$$

Indeed, the right hand side is contained in X^* . It suffices to show that $\varphi_j(X_I) \subset W'$ for every $j \in J$ (the symmetrical argument will show it for $\varphi_i(X_J)$, $i \in I$). Since $\pi_{abc}(\varphi_j(X_I)) \neq \{0\}$ if and only if exactly two of a, b, c are in I and either b or c are equal to j , this establishes our second claim.

Finally, we establish the equality in the lemma: let

$$\mathbf{v} \in \left(\bigcap_{i \in I} \varphi_i^{-1} \left(\langle \varphi_k(X_I) \mid k \in I \rangle \right) \right) \oplus \left(\bigcap_{j \in J} \varphi_j^{-1} \left(\langle \varphi_k(X_J) \mid k \in J \rangle \right) \right) \oplus V_{(I,J)}.$$

Then we can write $\mathbf{v} = \mathbf{x}_I \oplus \mathbf{x}_{(I,J)} \oplus \mathbf{x}_J$, where

$$\begin{aligned} \mathbf{x}_I &\in \bigcap_{i \in I} \varphi_i^{-1} \left(\langle \varphi_k(X_I) \mid k \in I \rangle \right), \\ \mathbf{x}_J &\in \bigcap_{j \in J} \varphi_j^{-1} \left(\langle \varphi_k(X_J) \mid k \in J \rangle \right), \\ \mathbf{x}_{(I,J)} &\in V_{(I,J)}. \end{aligned}$$

Since $V_{(I,J)} \subset X \subset X^{**}$, to show that $\mathbf{v} \in X^{**}$ it is enough to show that $\mathbf{x}_I$ and $\mathbf{x}_J$ are both in X^{**} . And indeed: notice that $\pi_{ab}(\mathbf{x}_I) \neq \mathbf{0}$ only if $a, b \in I$. Therefore, $\varphi_j(\mathbf{x}_I) \in W'$ for every $j \in J$. And by construction we have $\varphi_i(\mathbf{x}_I) \in X^*$ for every $i \in I$. Thus $\varphi_k(\mathbf{x}_I) \in X^*$ for all k . Symmetrically, $\varphi_k(\mathbf{x}_J) \in X^*$ for all k as well. Therefore, each of $\mathbf{x}_I$ and $\mathbf{x}_J$ lie in X^{**} , as desired.

For the converse inclusion, let $\mathbf{x} \in X^{**}$. We can write $\mathbf{x} = \mathbf{v}_I \oplus \mathbf{v}_J \oplus \mathbf{v}_{(I,J)}$, with $\mathbf{v}_I \in V_I$, $\mathbf{v}_J \in V_J$, and $\mathbf{v}_{(I,J)} \in V_{(I,J)}$. We must have $\varphi_k(\mathbf{v}) \in X^*$ for every k . If $k \in I$, then again we must have

$$\varphi_k(\mathbf{v}_I) \in \langle \varphi_i(X_I) \mid i \in I \rangle, \quad \varphi_k(\mathbf{v}_J), \varphi_k(\mathbf{v}_{(I,J)}) \in W'.$$

And if $k \in J$, then

$$\varphi_k(\mathbf{v}_J) \in \langle \varphi_j(X_J) \mid j \in J \rangle, \quad \varphi_k(\mathbf{v}_I), \varphi_k(\mathbf{v}_{(I,J)}) \in W'.$$

Therefore,

$$\begin{aligned} \mathbf{v}_I &\in \bigcap_{i \in I} \varphi_i^{-1}(\langle \varphi_k(X_I) \mid k \in I \rangle), \\ \mathbf{v}_J &\in \bigcap_{j \in J} \varphi_j^{-1}(\langle \varphi_k(X_J) \mid k \in J \rangle), \end{aligned}$$

as desired. This proves the equality.

The final clause of the lemma follows from the observation that X will be closed if and only if

$$X_I = \bigcap_{i \in I} \varphi_i^{-1}(\langle \varphi_k(X_I) \mid k \in I \rangle) \text{ and } X_J = \bigcap_{j \in J} \varphi_j^{-1}(\langle \varphi_k(X_J) \mid k \in J \rangle).$$

□

Notice that the subspace $V_{(I,J)}$, when interpreted as a subgroup of commutators we are making trivial, says that each x_i commutes with each x_j , where $i \in I$ and $j \in J$. From this we obtain the following:

Corollary 4.4. *Let G_1 and G_2 be two noncyclic p -groups of class at most two and exponent p . Then $G = G_1 \oplus G_2$ is capable if and only if each of G_1 and G_2 are capable.*

The corollary is clearly not true if we drop the “noncyclic” hypothesis, since a cyclic group of order p is not capable, but the direct sum of two cyclic groups of order p is capable. To understand why we must make the distinction, note that when $|I| = 1$, V_I is trivial so X_I is perforce $\{\varphi_i\}_{i \in I}$ -closed; but our developement always assumes $n > 1$. The symmetric situation happens when $|I| = n-1$. However, what we obtain in this instance is the following result, that is perhaps one of the most unexpected results from [6]:

Theorem 4.5 (Lemma 5.9 in [6]). *Let $n > 2$, and let X be a subspace of $V(n-1)$ and let $X' = X \oplus \langle v_{ni} \mid 1 \leq i \leq n-1 \rangle$ be a subspace of $V(n)$. Then X' is $\{\varphi_i\}_{i=1}^n$ -closed if and only if X is $\{\varphi_i\}_{i=1}^{n-1}$ -closed.*

The usefulness of this theorem becomes apparent when we translate it back into group theoretic terms:

Corollary 4.6. *Let K be a finite noncyclic group of exponent p and class 2, and let $G = K \oplus C_p$, where C_p is the cyclic group of order p . Then G is capable if and only if K is capable.*

Since every finite group of class at most two and exponent p may be written as $K \oplus C_p^r$ for some $r \geq 0$ and K satisfying $[K, K] = Z(K)$, we conclude that:

Corollary 4.7. *Let G be a p -group of class at most two and exponent p . Write $G = K \oplus C_p^r$, where K satisfies $[K, K] = Z(K)$. Then G is capable if and only if (i) K is nontrivial and capable, or (ii) K is trivial and $r \geq 2$.*

This allows us to ignore the “extra” elements in the center of G that do not result from commutators. Thus, for example, we can strengthen Theorem 3.5 by replacing the rank of G^{ab} by the rank of $G/Z(G)$. We then have the following:

Corollary 4.8. *Let G be a p -group of class two and exponent p , $\text{rank}(G/Z(G)) = n$, and $\text{rank}([G, G]) = \ell$. If $\ell > \binom{n}{2} - 2\sqrt{n}$, then G is capable.*

In [4], the authors established a minimum size for the commutator subgroup of a capable group G of class two and exponent p which satisfies $[G, G] = Z(G)$; namely, they proved:

Theorem 4.9 (Theorem 1 in [4]). *Suppose that G is a group of exponent p which satisfies $[G, G] = Z(G)$. If G is capable and $[G, G]$ is of rank ℓ , then $G/Z(G)$ is of rank at most $2\ell + \binom{\ell}{2}$.*

First, our development above now shows that if G is not cyclic, then Theorem 4.9 remains true if we drop the hypothesis that $[G, G] = Z(G)$. Second, note that Theorem 4.9 gives a *necessary* condition. If we let $n = \text{rank}(G/Z(G))$, as we did in Corollary 4.8, then Theorem 4.9 gives the inequality

$$\ell \geq \frac{\sqrt{9 + 8n} - 3}{2}$$

as a necessary condition for the capability of G . Combining it with Corollary 4.8, we have:

Theorem 4.10. *Let G be a noncyclic group of exponent p and class at most two. Let $\text{rank}(G/Z(G)) = n$ and $\text{rank}([G, G]) = \ell$. A necessary condition for the capability of G is that*

$$\ell \geq \frac{\sqrt{9 + 8n} - 3}{2}.$$

A sufficient condition for the capability of G is that

$$\ell > \binom{n}{2} - 2\sqrt{n}.$$

Although Theorem 4.5 focuses on the particular subspace that results when we take $I = \{1, 2, \dots, n-1\}$, in which case $V_{(j,i)}$ is none other than $\langle x_n \rangle^*$ (relative to $\{\psi_i\}_{i=1}^n$), there is nothing special about x_n (except that the theorem is easy to state). Because of the symmetry inherent in U , we can replace $\langle x_n \rangle^*$ with any other subspace of the form $\langle \mathbf{u} \rangle^*$, with $\langle \mathbf{u} \rangle \in \text{Gr}(1, U)$. It is easy to verify that in this case we will have $\langle \mathbf{u} \rangle \in \text{Gr}(n-1, V)$. Via a change of basis for U and the identifications $V = U \wedge U$, etc. noted at the end of Section 1, we obtain:

Theorem 4.11. *Fix $n > 2$, and let X' be a subspace of $V(n)$. If there exists $\langle \mathbf{u} \rangle \in \text{Gr}(1, U)$ such that $\langle \mathbf{u} \rangle^* \subset X'$, then there exists a subspace X of $V(n-1)$ of dimension $\dim(X') - n + 1$ such that X' is $\{\varphi_i\}_{i=1}^{n-1}$ -closed if and only if X is $\{\varphi_i\}_{i=1}^{n-1}$ -closed.*

5. THE CASE $n = 4$.

In this section we will settle the case $n \leq 4$. First, notice that for $n \leq 3$, Corollary 3.6 guarantees that all subspaces X will be closed. For $n = 4$, the same corollary settles all cases except $\dim(X) = 5$ and $\dim(X) = 6$; the latter case is just $X = V$, in which case $X^{**} = V = X$, so X is closed. Thus, we may restrict ourselves to $\dim(X) = 5$. Then the only two possibilities are $X^{**} = X$ and $X^{**} = V$. Since $X^{**} = V$ if and only if $X^* = W$, we have:

Proposition 5.1. *Let $n = 4$, and let X be a 5-dimensional subspace of V . Then X is closed if and only if X^* is a proper subspace of W .*

Since $\dim(W) = 20$, and in this case we have $\dim(W) = n \dim(V)$, the following observation becomes relevant:

Theorem 5.2. *Let $n > 1$ be an integer. Let X is a subspace of V , and assume that $\dim(X) \geq n$ and $\langle \mathbf{u} \rangle^* \subset X$ for some $\langle \mathbf{u} \rangle \in \text{Gr}(1, U)$. Then $\dim(X^*) < n \dim(X)$.*

Proof. By symmetry, it is enough to establish this result when $\mathbf{u} = x_1$. Then $\langle \mathbf{u} \rangle^*$ is generated by the vectors $v_{21}, v_{31}, \dots, v_{n1}$.

Let $\mathbf{v} \neq \mathbf{0}$ be a vector in X with $\Pi_1(\mathbf{v}) = \mathbf{0}$. Writing

$$\mathbf{v} = \sum_{2 \leq i < j \leq n} \alpha_{ji} v_{ji}$$

we have:

$$\begin{aligned} \varphi_1(\mathbf{v}) &= \sum_{2 \leq i < j \leq n} \alpha_{ji} (v_{j1i} - v_{i1j}) \\ &= \sum_{2 \leq i < j \leq n} (\varphi_i(\alpha_{ji} v_{j1}) - \varphi_j(\alpha_{ji} v_{i1})), \end{aligned}$$

so $\varphi_1(\mathbf{v}) \in \langle \varphi_2(X), \dots, \varphi_n(X) \rangle$. This shows that $\dim(X^*) < n \dim(X)$, as claimed. $\square$

For $n = 4$ and $\dim(X) = 5$, we claim that the converse holds.

Theorem 5.3. *Let $n = 4$, and let X be a 5-dimensional subspace of V . Then $\dim(X^*) < 20$ if and only if there exists a nonzero vector $\mathbf{u} \in U$ such that $\langle \mathbf{u} \rangle^* \subset X$.*

Proof. The if clause follows from Theorem 5.2. What we need to do is prove the converse. So assume that $\dim(X^*) < 20$. Let

$$Z_1 = \varphi_1(X) \cap \langle \varphi_2(X), \varphi_3(X), \varphi_4(X) \rangle \quad \text{and} \quad Z_2 = \varphi_2(X) \cap \langle \varphi_3(X), \varphi_4(X) \rangle.$$

At least one of Z_1 and Z_2 must be nontrivial.

If Z_2 is nontrivial, then it is dimension 1, and the only possibility is for v_{43} , v_{42} , and v_{32} to be elements of X . Let $\mathbf{v} = \alpha_{21} v_{21} + \alpha_{31} v_{31} + \alpha_{41} v_{41} \in X$ be a

nonzero element. This is possible since $\dim(X) = 5$. We claim that $\langle \mathbf{u} \rangle^* \subset X$, where $\mathbf{u} = \alpha_{21}x_2 + \alpha_{31}x_3 + \alpha_{41}x_4$. Indeed, we have:

$$\begin{aligned}\psi_1(\mathbf{u}) &= \alpha_{21}v_{21} + \alpha_{31}v_{31} + \alpha_{41}v_{41} = \mathbf{v} \in X; \\ \psi_2(\mathbf{u}) &= \alpha_{31}v_{32} + \alpha_{41}v_{42} \in X; \\ \psi_3(\mathbf{u}) &= -\alpha_{21}v_{32} + \alpha_{41}v_{43} \in X; \\ \psi_4(\mathbf{u}) &= -\alpha_{21}v_{42} - \alpha_{31}v_{43} \in X.\end{aligned}$$

Since each $\psi_i(\mathbf{u}) \in X$, it follows that $\langle \mathbf{u} \rangle^* \subset X$, as claimed.

If Z_1 is nontrivial, then there must be at least one nonzero vector in

$$\langle v_{32}, v_{42}, v_{43} \rangle \cap X$$

whose image under φ_1 lies in Z_1 . Say $\mathbf{v}_1 = \alpha v_{32} + \beta v_{42} + \gamma v_{43}$, with α, β, γ not all zero. Then X must also contain the following three vectors:

$$\begin{aligned}\mathbf{v}_2 &= \alpha v_{31} + \beta v_{41}, \\ \mathbf{v}_3 &= \alpha v_{21} - \gamma v_{41}, \\ \mathbf{v}_4 &= \beta v_{21} + \gamma v_{31},\end{aligned}$$

in order to obtain $\varphi_1(\mathbf{v}_1) = \varphi_2(\mathbf{v}_2) - \varphi_3(\mathbf{v}_3) - \varphi_4(\mathbf{v}_4)$.

However, $\mathbf{v}_2$, $\mathbf{v}_3$, and $\mathbf{v}_4$ are not linearly independent. For example, if $\gamma \neq 0$, then $\alpha \mathbf{v}_4 + \beta \mathbf{v}_3 = \alpha \gamma v_{31} - \beta \gamma v_{41} = \gamma(\alpha v_{31} + \beta v_{41}) = \gamma \mathbf{v}_2$, so $\mathbf{v}_2$ lies in $\langle \mathbf{v}_3, \mathbf{v}_4 \rangle$. Likewise, if $\beta \neq 0$ then $\mathbf{v}_3$ is a linear combination of $\mathbf{v}_2$ and $\mathbf{v}_4$; and if $\alpha \neq 0$ then $\mathbf{v}_4$ is a linear combination of $\mathbf{v}_2$ and $\mathbf{v}_3$.

If $\alpha \neq 0$, then taking a suitable scalar multiple of $\mathbf{v}$ we may assume $\alpha = 1$. So X contains the following three linearly independent vectors:

$$\begin{aligned}\mathbf{v}_1 &= v_{32} + \beta v_{42} + \gamma v_{43}, \\ \mathbf{v}_2 &= v_{31} + \beta v_{41}, \\ \mathbf{v}_3 &= v_{21} - \gamma v_{41},\end{aligned}$$

for some β, γ (possibly equal to zero). Since $\dim(X) = 5$, we must have at least two other linearly independent vectors. Adding suitable multiples of the $\mathbf{v}_j$, $j = 1, 2, 3$, we may assume that they both lie in $\langle v_{41}, v_{42}, v_{43} \rangle$.

If $v_{41} \in X$, then X contains v_{21} , v_{31} , and v_{41} , which means

$$\langle x_1 \rangle^* = \langle v_{21}, v_{31}, v_{41} \rangle \subset X.$$

If $v_{42}, v_{43} \in X$, then $v_{32} \in X$ as well, so Z_2 is nontrivial and as seen above there is some $\langle \mathbf{u} \rangle^* \subset X$. So we may assume that X has a basis consisting of $\mathbf{v}_1$, $\mathbf{v}_2$, and $\mathbf{v}_3$, together with two other vectors of the form:

$$\begin{aligned}\mathbf{v}_5 &= v_{41} + \delta v_{43}, \\ \mathbf{v}_6 &= v_{42} + \lambda v_{43}.\end{aligned}$$

In this case, let $\mathbf{u} = x_2 + \lambda x_3 + (\beta \lambda - \gamma)x_4$. Then we have:

$$\begin{aligned}\psi_1(\mathbf{u}) &= v_{21} + \lambda v_{31} + (\beta \lambda - \gamma)v_{41} = \mathbf{v}_3 + \lambda \mathbf{v}_2; \\ \psi_2(\mathbf{u}) &= \lambda v_{32} + (\beta \lambda - \gamma)v_{42} = \lambda \mathbf{v}_1 - \gamma \mathbf{v}_5; \\ \psi_3(\mathbf{u}) &= -v_{32} + (\beta \lambda - \gamma)v_{43} = -\mathbf{v}_1 + \beta \mathbf{v}_5; \\ \psi_4(\mathbf{u}) &= -v_{42} - \lambda v_{43} = -\mathbf{v}_5.\end{aligned}$$

Thus, $\langle \mathbf{u} \rangle^* \subset X$.

If $\alpha = 0$ and $\beta \neq 0$, then we may assume $\beta = 1$, so now the following vectors lie in X :

$$\begin{aligned}\mathbf{v}_1 &= v_{42} + \gamma v_{43}, \\ \mathbf{v}_2 &= v_{41}, \\ \mathbf{v}_4 &= v_{21} + \gamma v_{31}.\end{aligned}$$

Two other vectors, which we may assume lie in $\langle v_{31}, v_{32}, v_{43} \rangle$, will form a basis for X . If $v_{43} \in X$, then $\langle x_4 \rangle^* = \langle v_{31}, v_{32}, v_{43} \rangle \subset X$, and we are done. Otherwise, we may assume that

$$\begin{aligned}\mathbf{v}_5 &= v_{31} + \delta v_{43}, \\ \mathbf{v}_6 &= v_{32} + \lambda v_{43},\end{aligned}$$

lie in X , for some δ, λ . In this case, let $\mathbf{u} = x_2 + \gamma x_3 - \lambda x_4$, and we obtain

$$\begin{aligned}\psi_1(\mathbf{u}) &= \mathbf{v}_4 - \lambda \mathbf{v}_2, \\ \psi_2(\mathbf{u}) &= \gamma \mathbf{v}_6 - \lambda \mathbf{v}_1, \\ \psi_3(\mathbf{u}) &= -\mathbf{v}_6, \\ \psi_4(\mathbf{u}) &= -\mathbf{v}_1.\end{aligned}$$

Finally, if $\alpha = \beta = 0$, we may assume $\gamma = 1$, so now we have the following linearly independent vectors in X :

$$\begin{aligned}\mathbf{v}_1 &= v_{43}, \\ \mathbf{v}_3 &= -v_{41}, \\ \mathbf{v}_4 &= v_{31},\end{aligned}$$

plus two more vectors to form a basis, these latter vectors taken from $\langle v_{21}, v_{32}, v_{42} \rangle$. If $v_{21} \in X$, then $\langle x_1 \rangle^* = \langle v_{21}, v_{31}, v_{41} \rangle \subset X$, and again we are done. Otherwise, we may take

$$\begin{aligned}\mathbf{v}_5 &= v_{32} + \delta v_{21}, \\ \mathbf{v}_6 &= v_{42} + \lambda v_{21},\end{aligned}$$

as two other vectors in X . This time, letting $\mathbf{u} = -\delta x_1 + x_3$ gives $\langle \mathbf{u} \rangle^* \subset X$, as desired. $\square$

Corollary 5.4. *Let $n = 4$, and let X be a 5-dimensional subspace of V . Then X is closed if and only if there exists a nonzero vector $\mathbf{u} \in U$ such that $\langle \mathbf{u} \rangle^* \subset X$.*

This result may seem a bit artificial when put in this guise. However, translating it back to group theory reveals what is going on. Recall that the vectors in U correspond to elements of G^{ab} , and if $\langle \mathbf{u} \rangle^* \subset X$, that means that the corresponding element of G is central (since commutators are central in G , the ambiguity in the statement “the corresponding element of G ” is immaterial). So, in group theoretic language, Corollary 5.4 states:

Theorem 5.5. *Let G be a group of class two and exponent p , p and odd prime. If G is of order p^5 and $[G, G]$ is cyclic of order p , then G is capable if and only if $[G, G] \neq Z(G)$. That is, in this case, G is capable if and only if G is not extra-special.*

Recall that we say a group G is n -generated to mean that it can be generated by n elements, though it may in fact need fewer. We say G is *minimally n -generated* to mean that G can be generated by n elements, but cannot be generated by k -elements for any $k < n$.

Let G be any 4-generated group of exponent p and class at most 2, p an odd prime. If G is cyclic and nontrivial, then it is neither capable nor extra-special. If G is minimally 2- or 3-generated, then it is capable. If G is minimally 4-generated, then it is capable in all cases except when G is extra-special of order p^5 . Therefore, we have established the following:

Theorem 5.6. *Let G be a 4-generated p -group of class at most two and exponent p . Then G satisfies one and only one of the following:*

- (i) G is capable; or
- (ii) G is cyclic and nontrivial; or
- (iii) G is extra-special of order p^5 and exponent p .

Remark 5.7. It may be profitable to consider the following: each $\langle \mathbf{u} \rangle \in Gr(1, U)$ determines, via $\langle \mathbf{u} \rangle^*$, an $(n - 1)$ -dimensional subspace of V . For every $k \geq n - 1$, this subspace determines a well-known subset (in fact, a subvariety) of $Gr(k, V)$, known as *Schubert cells* or *Schubert cycles*. For the case $n = 4$, the above theorem states that the $X \in Gr(5, V)$ that are closed are exactly the ones that are in some Schubert cycle determined as $\langle \mathbf{u} \rangle$ ranges over all elements of $Gr(1, U)$.

REFERENCES

- [1] Michael R. Bacon and Luise-Charlotte Kappe, *On capable p -groups of nilpotency class two*, Illinois J. Math. **47** (2003), 49–62. MR **2004j**:20036
- [2] Reinhold Baer, *Groups with preassigned central and central quotient group*, Trans. Amer. Math. Soc. **44** (1938), 387–412.
- [3] F. Rudolf Beyl, Ulrich Felgner, and Peter Schmid, *On groups occurring as central factor groups*, J. Algebra **61** (1979), 161–177. MR **81i**:20034
- [4] Hermann Heineken and Daniela Nikolova, *Class two nilpotent capable groups*, Bull. Austral. Math. Soc. **54** (1996), 347–352. MR **97m**:20043
- [5] Arturo Magidin, *Capability of nilpotent products of cyclic groups*, arXiv:math.GR/0403188, to appear in J. Group Theory.
- [6] ———, *Capable groups of prime exponent and class two*, arXiv:math.GR/0401423, Submitted.
- [7] ———, *Capable two-generator 2-groups of class two*, arXiv:math.GR/0404499, Submitted.
- [8] David McKinnon, Personal communication.

DEPT. MATHEMATICAL SCIENCES, THE UNIVERSITY OF MONTANA, MISSOULA MT 59812-0864
E-mail address: `magidin@member.ams.org`